

Stoichiometric and temporal requirements of Oct4, Sox2, Klf4, and c-Myc expression for efficient human iPSC induction and differentiation

Eirini P. Papapetrou^{a,b}, Mark J. Tomishima^{c,d}, Stuart M. Chambers^c, Yvonne Mica^e, Evan Reed^{a,b}, Jayanthi Menon^{a,f}, Viviane Tabar^{a,f}, Qianxing Mo^g, Lorenz Studer^{a,c}, and Michel Sadelain^{a,b,1}

^aCenter for Cell Engineering, ^bMolecular Pharmacology and Chemistry and ^cDevelopmental Biology Programs, ^dSKI Stem Cell Research Facility, ^eGerstner Sloan-Kettering Graduate School, and Departments of ^fNeurosurgery and ^gEpidemiology and Biostatistics, Memorial Sloan-Kettering Cancer Center, New York, NY, 10065

Communicated by Mark Ptashne, Memorial Sloan-Kettering Cancer Center, New York, NY, May 1, 2009 (received for review April 8, 2009)

Human-induced pluripotent stem cells (hiPSCs) are generated from somatic cells by ectopic expression of the 4 reprogramming factors (RFs) Oct-4, Sox2, Klf4, and c-Myc. To better define the stoichiometric requirements and dynamic expression patterns required for successful hiPSC induction, we generated 4 bicistronic lentiviral vectors encoding the 4 RFs co-expressed with discernable fluorescent proteins. Using this system, we define the optimal stoichiometry of RF expression to be highly sensitive to Oct4 dosage, and we demonstrate the impact that variations in the relative ratios of RF expression exert on the efficiency of hiPSC induction. Monitoring of expression of each individual RF in single cells during the course of reprogramming revealed that vector silencing follows acquisition of pluripotent cell markers. Pronounced lentiviral vector silencing was a characteristic of successfully reprogrammed hiPSC clones, but lack of complete silencing did not hinder hiPSC induction, maintenance, or directed differentiation. The vector system described here presents a powerful tool for mechanistic studies of reprogramming and the optimization of hiPSC generation.

fluorescent proteins | lentiviral vectors | silencing | stoichiometry

Reprogramming of human fibroblasts to a pluripotent embryonic stem cell (ESC)-like state has recently been achieved through retroviral-mediated gene transfer of the 4 transcription factors Oct-4, Sox-2, Klf-4, and c-Myc (1–3). This combination of factors emerged from an initial screen in mouse fibroblasts based on co-transduction of 24 candidate genes (4). The generation of iPSCs with this method has now been reported from mouse and human somatic cells using various types of vectors, including gamma-retroviral, constitutive, or doxycycline (DOX)-inducible lentiviral and adenoviral vectors, as well as plasmid and transposon/transposase transfection systems (1, 2, 5–12). Nonetheless, the stoichiometric and temporal requirements of factor expression during hiPSC induction, maintenance, and differentiation remain poorly defined.

Direct reprogramming is a slow and inefficient process, with estimated efficiencies in human cells ranging from 0.02% to 0.002% (1, 2, 5). Many on-going efforts aim to identify genetic or chemical factors that enhance iPSC generation (13–16). A problem faced by these investigations is the lack of a consistent way of reporting reprogramming efficiency, since effects on factor delivery cannot be separated from genuine effects on reprogramming efficiency. Furthermore, the lack of proper assessment of factor delivery and expression obscures the comparison of reprogramming frequencies across different studies. The low efficiency of direct reprogramming may, at least in part, be accounted for by the requirement for a stringent stoichiometry of reprogramming factor expression permissive for successful reprogramming.

Several studies have reported silencing of the vector-encoded reprogramming factors in iPSCs generated with gamma-retroviral vectors (1, 17, 18). The silencing of lentiviral vectors in iPSCs is less known (2, 19), and therefore their suitability for reprogramming

purposes remains controversial (20). Furthermore, the significance of vector silencing for reprogramming and its impact on differentiation is unclear. Although it has been proposed that silencing of factor expression is required for successful reprogramming and multilineage differentiation (3, 21, 22), it remains unclear whether vector silencing constitutes a requirement or an epiphenomenon of the reprogramming process.

Here we present a vector system for iPSC induction that allows for simultaneous real-time tracking of expression of the 4 individual transgenes in single cells during hiPSC induction, maintenance and directed differentiation. Using this system, we show that expression of the 4 RFs at an optimal stoichiometry is critical for efficient reprogramming, and we study in detail the kinetics of silencing or the 4 transgenes during the reprogramming process.

Results

Reprogramming of Human Fibroblasts with Bicistronic Vectors Coexpressing Each Reprogramming Factor with a Distinct Fluorescent Protein. We constructed lentiviral vectors that co-express each of the 4 RFs, Oct4, Sox2, Klf4, and c-Myc together with a fluorescent protein linked by a 2A peptide (Fig. 1A). We selected a combination of 4 fluorescent proteins, vGFP (violet light excited-green fluorescent protein), mCitrine, mCherry, and mCerulean (23, 24), which can be separated by fluorescence microscopy and flow cytometry, thus enabling parallel monitoring of each reprogramming factor expression in real time (Fig. 1B and see below). Immunoblots of cells transduced with these vectors revealed correct processing of the 2A-linked gene products (Fig. S1 A–D). All 4 vectors were found to yield similar titers of $1.5\text{--}2 \times 10^6$ TU/mL. By co-transduction of human fetal fibroblasts (MRC-5) with these 4 vectors hiPSC lines were generated that expressed pluripotency markers (Fig. S2A) and could be directed to differentiate into derivatives of all 3 germ layers (Fig. S2B). Microarray-based global gene expression analysis revealed gene expression patterns highly similar to hESCs, but not to the parental human fibroblasts (Fig. S2C and Table S1). Furthermore, these hiPSC lines induced teratomas when s.c. injected into immunocompromised mice (Fig. 1C).

Requirements of Factor Expression for Efficient Reprogramming. We first took advantage of this traceable vector system to estimate what fraction of human fibroblasts co-expressing all 4 factors successfully reprogram (Fig. S2D). We co-transduced human fetal fibroblasts (MRC-5) with the 4 vectors at equal multiplicity of infection (MOI)

Author contributions: E.P.P., M.J.T., L.S., and M.S. designed research; E.P.P., M.J.T., S.M.C., Y.M., E.R., J.M., and V.T. performed research; E.P.P., Q.M., and M.S. analyzed data; and E.P.P. and M.S. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: m-sadelain@ski.mskcc.org.

This article contains supporting information online at www.pnas.org/cgi/content/full/0904825106/DCSupplemental.




Fig. 1. hiPSCs derived from bicistronic vectors co-expressing each RF linked to a fluorescent protein. (A) Schematic representation of the vectors used in this study. LTR, long terminal repeat. (B) hiPSC colony at day 11 after transduction. (C) Hematoxylin and eosin staining of histological sections of a teratoma derived from line iP5-27. From *Left to Right*: Low power image demonstrating areas of heterogeneous differentiation: neuroectoderm (black arrow), smooth muscle (blank arrow), and primitive myxoid tissue (arrowhead), 4 \times . Pigmented epithelial tissue compatible with retinal neuroectoderm, 40 \times . Intestinal like epithelium including goblet cells (endoderm), 40 \times . Smooth muscle tissue, 20 \times . Inset demonstrates a high power image of immature mesenchymal tissue, potentially cartilage (40 \times).

and determined the percentage of quadruple transduced cells by flow cytometry 5 days after transduction. To calculate the frequency of fully reprogrammed cells, we performed immunostaining for Tra-1-81, as described in the *Methods*, and macroscopically enumerated positive colonies on day 20 after transduction. Thus, through direct quantification of quadruple transduced cells coupled with a clonal readout of hiPSC derivation, we were able to accurately estimate the reprogramming frequency to be 0.4–1% of de novo quadruple transduced fetal human fibroblasts (Table S2).

We then sought to investigate the effect of the absolute levels of expression of the 4 vector-encoded RFs on efficiency of direct reprogramming. Transduction of human fibroblasts at increasing MOI results in concomitant increase in the percentage of quadruple transduced cells, as well as in the level of expression (Fig. 2*A* and *B*). Importantly, titration of the MOI of each vector results in linear titration of the mean fluorescence intensity (MFI) of the corresponding fluorescent protein (Fig. 2*B*). Our vector design also ensures that titrated expression of each fluorescent protein corresponds to titrated levels of expression of each linked RF (Fig. S1*E*). Increasing the levels of expression of the 4 transgenes simultaneously, as exemplified in Fig. 2*B*, had no effect in reprogramming efficiency of quadruple transduced cells. This result was corroborated by fluorescence-activated cell sorting (FACS) experiments that showed that the reprogramming efficiency of quadruple transduced human fibroblasts expressing all 4 transgenes at similarly high or low levels was the same (Fig. S3).

We therefore sought to investigate the effect that stoichiometric deviations in factor expression impose in the efficiency of reprogramming. We hypothesized that if differences in ratios of RF expression affect reprogramming efficiency in single cells, we could observe similar trends in bulk populations of quadruple transduced cells. Starting from a vector proportion of 1:1:1:1, we altered the stoichiometric ratio by varying the amount of 1 vector at a time while keeping the other 3 constant, as detailed in Fig. 2*C*. Flow cytometric analysis of factor expression 5 days after transduction was used to identify groups in which expression levels of any 3 factors (provided in constant amounts) maintained the same MFI and those were selected for comparison across permutations of expression of the fourth factor (Fig. S4*A* and *B*). Reprogramming efficiency was calculated by enumeration of Tra-1-81+ colonies as above (Fig. S4*C*). We again did not observe correlation of reprogramming efficiency with the absolute expression levels of any of

the 4 factors. In contrast, changes in the relative ratios of expression mediated significant effects on the efficiency of reprogramming. Increasing relative Oct4 expression resulted in enhanced reprogramming efficiency, whereas increasing the relative ratio of either Sox2, Klf4, or c-Myc consistently decreased efficiency of hiPSC colony generation by more than 5-fold (Fig. 2*C* *Upper*). In contrast, relative decrease of Sox2, Klf4 or c-Myc showed little effect, while relative decrease of Oct4 was detrimental (Fig. 2*C* *Lower*). These data demonstrate that a stoichiometry of equal parts of all 4 vectors is highly effective, resulting in reprogramming efficiency of de novo quadruple transduced fibroblasts similar to that observed in the maximally efficient secondary system (5, 7). Most deviations from this 1:1:1:1 stoichiometry have unfavorable effects to the efficiency of reprogramming, with the exception of a relative Oct4 increase. The amount of Oct4 expression has been shown to be critical in ES cells and its up- or down- regulation both drastically alter the pluripotent cell phenotype (25). We therefore examined in more detail the effect of relative Oct4 expression levels in direct reprogramming. Although a 3-fold relative increase was favorable, further increases were detrimental (Fig. 2*D*). Similar titrations of relative expression of the remaining 3 factors over a wider range did not reveal any additional effect than the trends shown in Fig. 2*C*. Based on these results, the optimal RF stoichiometry consists of a combination of equal amounts of Sox2, Klf4, and c-Myc with a 3-fold excess of Oct4.

Pronounced Transgene Silencing Is a Hallmark of Successful Reprogramming but Residual Factor Expression Does Not Hinder Early Lineage Specification. To study vector silencing, we established a panel of successfully reprogrammed clones on the basis of hESC-like morphology, Tra-1-81 expression and HLA-ABC downregulation to levels similar to hESCs ($n = 38$) and a panel of clones that formed colonies with non-hES morphology, did not express pluripotency markers and retained high HLA-ABC expression ($n = 30$). Ectopic expression of all 4 RFs was markedly lower in the panel of hiPSC lines than in lines that had not successfully undergone reprogramming to hiPSC state (Figs. 3 and S5*A*).

Despite showing pronounced vector silencing, a number of hiPSC lines were found not to have completely silenced factor expression. Some hiPSC lines maintained expression of 1 or more factors (at levels 20-fold lower than in transduced fibroblasts and in not fully reprogrammed clones). It has been suggested that some




Fig. 4. Differentiation of hiPSC clones with incomplete RF silencing. *Left*, flow cytometry analysis of hiPSC clones 72, 71, 62 and 50. *Right*, in vitro differentiation of the same clones in endoderm, mesoderm and ectoderm, followed by staining for Sox17, Brachyury, and Pax6, respectively.

acquisition of several criteria of pluripotency, such as morphology, growth, and pluripotency marker expression.

We further hypothesized that, if silencing were a requirement for reprogramming, it would be selective for the reprogramming

vectors. To test this, we examined the state of silencing in fully reprogrammed (Tra-1-81+) versus partially reprogrammed (Tra-1-81-) clones generated from MRC-5 fibroblasts co-transduced with the 4 bicistronic vectors and an additional fifth




Fig. 5. Real-time simultaneous monitoring of vector-encoded RF expression and pluripotency marker induction. (A) Vector expression in Tra-1-81+ versus HLA-ABC^{high} cells on day 12 (Left) and 20 (Right) after transduction. (B) Vectorsilencing in Tra-1-81+ cells over time. (Day 0 denotes time of transduction.) Fold difference was estimated as ratio of MFI in the Tra-1-81+ cells to the MFI in HLA-ABC^{high} cells. (C) Appearance of pluripotency marker Tra-1-81 and downregulation of HLA-ABC during reprogramming. Shown is 1 of 3 independent time course experiments.

secondary antibodies conjugated to Alexa488 or Alexa568 (Invitrogen) were used for detection. Nuclei were detected with Hoechst 33258.

For Tra-1-81 immunostaining, duplicate plates were fixed with 4% para-formaldehyde and incubated with an anti-Tra-1-81 antibody (Chemicon), followed by incubation with a horseradish peroxidase-linked anti-mouse IgM secondary antibody (Invitrogen). Staining of positive colonies was achieved by addition of chromogenic substrate 4-chloro-1-naphthol (Sigma). Macroscopic enumeration of positive colonies was done by 2 independent reviewers in a blinded fashion.

Gene Expression Profiling. Total RNA was isolated with RNeasy kit (QIAGEN). Samples were processed as independent triplicates. Whole genome gene expression analysis was performed on Illumina BeadArrays at the MSKCC microarray facility. Software R (<http://www.r-project.org/>) was used to perform all statistical computations. Moderated 2-sample t-test implemented in LIMMA package was used to examine whether genes were differentially expressed. Storey's *q*-value that controls positive false discovery rate (FDR) was used to correct for multiple testing for each contrast of interest. *Q*-values less than 0.05 were considered statistically significant.

Teratoma Formation. Undifferentiated hiPSCs were suspended at 1×10^7 cells/mL in DMEM containing 10% FBS. One million cells were injected s.c. in the flank of adult (3-month-old) NOD/SCID mice (Jackson Laboratory). Eight weeks later, the tumor was surgically dissected, fixed in 4% formaldehyde and embedded in OCT. Sections were cut at 7- μ m thickness and stained with hematoxylin and eosin. All animal experiments were conducted in accordance with protocols approved by MSKCC Institutional Animal Care and Use Com-

mittee (IACUC) and following National Institutes of Health guidelines for animal welfare.

Flow Cytometry. Cells were dissociated with accutase, stained with Alexa Fluor (AF) 647-conjugated anti-Tra-1-81 and PE-Cy5-conjugated anti-HLA-ABC or AF 647-conjugated anti-CXCR4 antibodies (BD Biosciences) and analyzed in a LSRII cytometer (BD Biosciences). Analysis was performed with the FlowJo software (version 8.8.4; Tree Star). Cell sorting was performed on a MoFlo cell sorter (DakoCytomation).

In Vitro Differentiation. Endoderm and mesoderm were induced as previously described (37, 38). Briefly, for endoderm differentiation hiPSCs were passaged onto MEFs and expanded for 2–3 days before media was switched to endoderm induction medium (RPMI, 0.5% FBS, 2 mM L-glutamine, and 100 ng/mL activin A). For mesoderm differentiation hiPSCs were passaged onto MEFs and expanded for 6 days before media was switched to mesoderm induction medium (DMEM/F12, 10% FBS, 2 mM L-glutamine, 0.1 mM nonessential amino acids, and 0.1 mM beta-mercaptoethanol). In both protocols, the medium was replaced every other day and the cells were assayed on day 6. Neural induction was done as previously described (39).

Statistical Analysis. Linear regression analysis was performed using Prism software (version 5.0a; GraphPad).

ACKNOWLEDGMENTS. The authors would like to thank Kim Ping for excellent technical assistance, Ronald Grosse for assistance in histological analysis, Christian Carson and Jurg Rohrer (BD Biosciences) and Jan Hendrikx for cell sorting. This work was supported by the Starr Foundation (Tri-Institutional Stem Cell Initiative, Tri-SCI-018) and New York State Stem Cell Science, NYSTEM (N08T-060).

1. Takahashi K, et al. (2007) Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 131:861–872.
2. Yu J, et al. (2007) Induced pluripotent stem cell lines derived from human somatic cells. *Science* 318:1917–1920.
3. Park IH, et al. (2008) Reprogramming of human somatic cells to pluripotency with defined factors. *Nature* 451:141–146.
4. Takahashi K, Yamanaka S (2006) Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 126:663–676.
5. Maherali N, et al. (2008) A high-efficiency system for the generation and study of human induced pluripotent stem cells. *Cell Stem Cell* 3:340–345.
6. Wernig M, et al. (2008) A drug-inducible transgenic system for direct reprogramming of multiple somatic cell types. *Nat Biotechnol* 26:916–924.
7. Hockemeyer D, et al. (2008) A drug-inducible system for direct reprogramming of human somatic cells to pluripotency. *Cell Stem Cell* 3:346–353.
8. Stadtfeld M, Nagaya M, Utikal J, Weir G, Hochedlinger K (2008) Induced pluripotent stem cells generated without viral integration. *Science* 322:945–949.
9. Okita K, Nakagawa M, Hyenjong H, Ichisaka T, Yamanaka S (2008) Generation of mouse induced pluripotent stem cells without viral vectors. *Science* 322:949–953.
10. Kaji K, et al. (2009) Virus-free induction of pluripotency and subsequent excision of reprogramming factors. *Nature* 458:771–775.
11. Woltjen K, et al. (2009) piggyBac transposition reprograms fibroblasts to induced pluripotent stem cells. *Nature* 458:766–770.
12. Yu J, et al. (2009) Human induced pluripotent stem cells free of vector and transgene sequences. *Science* 324:797–801.
13. Mikkelsen TS, et al. (2008) Dissecting direct reprogramming through integrative genomic analysis. *Nature* 454:49–55.
14. Shi Y, et al. (2008) Induction of pluripotent stem cells from mouse embryonic fibroblasts by Oct4 and Klf4 with small-molecule compounds. *Cell Stem Cell* 3:568–574.
15. Zhao Y, et al. (2008) Two supporting factors greatly improve the efficiency of human iPS generation. *Cell Stem Cell* 3:475–479.
16. Huangfu D, et al. (2008) Induction of pluripotent stem cells by defined factors is greatly improved by small-molecule compounds. *Nat Biotechnol* 26:795–797.
17. Brambrink T, et al. (2008) Sequential expression of pluripotency markers during direct reprogramming of mouse somatic cells. *Cell Stem Cell* 2:151–159.
18. Stadtfeld M, Maherali N, Breault DT, Hochedlinger K (2008) Defining molecular cornerstones during fibroblast to iPS cell reprogramming in mouse. *Cell Stem Cell* 2:230–240.
19. Blueloch R, Venere M, Yen J, Ramalho-Santos M (2007) Generation of induced pluripotent stem cells in the absence of drug selection. *Cell Stem Cell* 1:245–247.
20. Maherali N, Hochedlinger K (2008) Guidelines and techniques for the generation of induced pluripotent stem cells. *Cell Stem Cell* 3:595–605.
21. Jaenisch R, Young R (2008) Stem cells, the molecular circuitry of pluripotency and nuclear reprogramming. *Cell* 132:567–582.
22. Wernig M, et al. (2007) In vitro reprogramming of fibroblasts into a pluripotent ES-cell-like state. *Nature* 448:318–324.
23. Anderson MT, et al. (1998) Pairs of violet-light-excited fluorochromes for flow cytometric analysis. *Cytometry* 33:435–444.
24. Shaner NC, et al. (2004) Improved monomeric red, orange, and yellow fluorescent proteins derived from *Drosophila* sp. red fluorescent protein. *Nat Biotechnol* 22:1567–1572.
25. Niwa H, Miyazaki J, Smith AG (2000) Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nat Genet* 24:372–376.
26. Teich NM, Weiss RA, Martin GR, Lowy DR (1977) Virus infection of murine teratocarcinoma stem cell lines. *Cell* 12:973–982.
27. Hotta A, Ellis J (2008) Retroviral vector silencing during iPS cell induction: An epigenetic beacon that signals distinct pluripotent states. *J Cell Biochem* 105:940–948.
28. Silva J, et al. (2008) Promotion of reprogramming to ground state pluripotency by signal inhibition. *PLoS Biol* 6:e253.
29. Eminli S, Utikal J, Arnold K, Jaenisch R, Hochedlinger K (2008) Reprogramming of neural progenitor cells into induced pluripotent stem cells in the absence of exogenous Sox2 expression. *Stem Cells* 26:2467–2474.
30. Sridharan R, et al. (2009) Role of the murine reprogramming factors in the induction of pluripotency. *Cell* 136:364–377.
31. Sommer CA, et al. (2008) Induced pluripotent stem cell generation using a single lentiviral stem cell cassette. *Stem Cells* 27:543–549.
32. Carey BW, et al. (2008) Reprogramming of murine and human somatic cells using a single polycistronic vector. *Proc Natl Acad Sci USA* 106:157–162.
33. Soldner F, et al. (2009) Parkinson's disease patient-derived induced pluripotent stem cells free of viral reprogramming factors. *Cell* 136:964–977.
34. Mali P, et al. (2008) Improved efficiency and pace of generating induced pluripotent stem cells from human adult and fetal fibroblasts. *Stem Cells* 26:1998–2005.
35. Szymczak AL, et al. (2004) Correction of multi-gene deficiency in vivo using a single 'self-cleaving' 2A peptide-based retroviral vector. *Nat Biotechnol* 22:589–594.
36. Papapetrou EP, Kovalovsky D, Beloeil L, Sant'angelo D, Sadelain M (2009) Harnessing endogenous miR-181a to segregate transgenic antigen receptor expression in developing versus post-thymic T cells in murine hematopoietic chimeras. *J Clin Invest* 119:157–168.
37. Watanabe K, et al. (2007) A ROCK inhibitor permits survival of dissociated human embryonic stem cells. *Nat Biotechnol* 25:681–686.
38. D'Amour KA, et al. (2005) Efficient differentiation of human embryonic stem cells to definitive endoderm. *Nat Biotechnol* 23:1534–1541.
39. Chambers SM, et al. (2009) Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nat Biotechnol* 27:275–280.